



Diabetes in India: what is different?

Viral N. Shah^{a,*} and Viswanathan Mohan^{b,*}

Purpose of review

The purpose of this study was to review the epidemiology and pathogenesis of diabetes in Asian Indians with a particular focus on 'Asian Indian type 2 diabetes phenotype'.

Recent findings

The prevalence of diabetes is rapidly increasing among Asian Indians, particularly in the past two decades. The diabetes rates in urban India now exceed that seen in Indians migrated to developed nations. Urbanization, changes from traditional healthier diets to high-refined carbohydrate intake, and sedentary lifestyle have contributed to this steep increase in the prevalence of diabetes in India. Type 2 diabetes among Asian Indians is characterized by onset at a younger age, greater abdominal obesity despite relatively lower BMI, greater insulin resistance, and early decline in beta cell function. Asian Indians are also at a higher risk for premature coronary artery disease.

Summary

The clinical profile of type 2 diabetes in Asian Indians differs from Caucasians with higher central obesity, increased inflammatory markers such as high sensitive C-reactive protein, greater insulin resistance, early loss of beta cell function, and a higher risk of coronary artery disease. Mechanistic studies are needed to characterize the pathophysiology of the Asian Indian phenotype.

Keywords

Asian Indian phenotype, Asian Indians, epidemiology, prevalence, south Asian phenotype, south Asians, type 2 diabetes

INTRODUCTION

Diabetes was known to Indian physicians even before 1500 BC [1]. Several ancient texts including the well known Ayurvedic texts (Charak Samhita, 1000 BC and Susruta, 600 BC) gave detailed descriptions of its symptoms and classified diabetes into lean or hereditary and obese or food-induced diabetes, which is not very dissimilar to the present-day classification of diabetes as type 1 (T1D) and type 2 diabetes (T2D) [2].

There is circumstantial evidence from ancient texts and Indian sculptures and paintings depicting large bellied individuals to suggest that obesity and T2D were not uncommon in ancient India [2,3]. Although the exact prevalence of diabetes during the precolonial and colonial periods is not known, it is believed that the repeated invasion by the Arabs and Mughals followed by British colonization leads to impoverishment that might be one of the reasons for the lower prevalence of diabetes during postcolonial independent India [2,3]. Since 1991, socioeconomic conditions in India have dramatically improved, resulting in a rapid increase in the prevalence of T2D. Thus, the increase in prevalence of diabetes in India is one of

the major health concerns posing a great economic burden on developing India.

We have reviewed the recent research on epidemiology of diabetes, factors contributed to a rise in diabetes, and characteristics of T2D among Asian Indians.

Epidemiology of diabetes in India

Since the 1960s, a number of studies have estimated the prevalence of T2D in various regions of India. Only multicenter studies are, however, discussed

^aBarbara Davis Center for Diabetes, University of Colorado Anschutz Campus, Aurora, Colorado, USA and ^bMadras Diabetes Research Foundation and Dr Mohan's Diabetes Specialties Centre, Chennai, India

Correspondence to Dr Viral N. Shah, MD, Assistant Professor of Medicine and Pediatrics, Adult Clinic, Barbara Davis Center for Diabetes, University of Colorado, 1775 Aurora Ct, Mail stop A140, Aurora, CO 80045, USA. Tel: +1 303 724 8186; fax: +1 303 724 6784; e-mail: viral.shah@ucdenver.edu

*Viral N. Shah and Viswanathan Mohan contributed equally to the writing of this article.

Curr Opin Endocrinol Diabetes Obes 2015, 22:283–289

DOI:10.1097/MED.000000000000166

KEY POINTS

- The prevalence of T2D in both urban and rural India has increased 10-fold in the last two decades.
- Urbanization, change from traditional healthier diets to those high in refined carbohydrates and fat, and decreased physical activity have contributed to a steep increase in the prevalence of diabetes in India.
- The epidemiology of T2D is changing with the shift of youth to poor urban and rural areas.
- The 'Asian Indian phenotype' is characterized by the onset of T2D at a younger age, greater abdominal obesity despite relatively lower BMI, greater insulin resistance, and early decline in beta cell function.

here. The first national study on the prevalence of T2D in India by the Indian Council of Medical Research (ICMR) was conducted between 1972 and 1975 found the prevalence of T2D in 2.1% of the urban population and 1.5% of the rural population [4]. The 1980s saw a slow, but definite, rise in the diabetes epidemic in both urban and rural India [5,6], and therefore, ICMR conducted a second survey of the prevalence of diabetes. It was reported as 4.1% among urban and 1.5% in rural areas that was almost double compared with the first national survey, especially in urban India [7]. In 2001, a population-based study – National Urban Diabetes Survey – from six large Indian metropolitan cities reported that the age standardized prevalence of diabetes was 12.4% with large variations between different cities [8]. In 2004, the Prevalence of Diabetes in India Study was carried out in 108 centers (49 urban and 59 rural) in different parts of India but excluded metro cities. This study showed a prevalence of diabetes of 5.6% and 2.7% among urban and rural areas, respectively [9]. Another survey, conducted in 2008 in an industrial cohort from five sites in India, showed the prevalence of diabetes in this population was 10.1% [10].

The ICMR-INDIA DIABetes (INDIAB) study is the largest nationwide study on the prevalence of diabetes and is studying all 29 Indian states, two union territories, and the National Capital Region of Delhi in a phased manner. Phase 1 of the ICMR-INDIAB study was carried out in four Indian states, namely, Tamil Nadu (South), Chandigarh (Union Territory, North), Maharashtra (West), and Jharkhand (East). This study reported the prevalence of diabetes (both known and newly diagnosed together) varied from 5 to 13% in different parts of India. Based on these, the overall number of

people with diabetes and prediabetes in year 2011 was estimated at 62.4 million and 77.2 million, respectively [11].

Thus, the number of people with diabetes in India has increased from 19 million in 1995 [12] to 32 million in 2000 [13] to 62.4 million in 2011 [11] to 66.8 million in 2014 [14]. The various epidemiological studies conducted over this period of time show at least a 10-fold rise in the prevalence of diabetes in the rural areas is quickly catching up with the urban areas in the prevalence of diabetes in India. The trend of rising prevalence of diabetes over this period of time from various multicenter national studies is shown in Fig. 1.

Rising prevalence of early onset type 2 diabetes among Indians

One of the most worrisome aspects of T2D in Indians is the diabetes onset at a young age. Studies have shown higher prevalence of T2D among young adult Indians [11,15,16]. In a recent clinic-based study from the southern part of India, the prevalence of diabetes in the young (diagnosed ≤ 25 years of age) rose significantly from 0.55% in 1992 to 2.5% in 2009, and this is largely because of the increasing number of T2D, whereas T1D has remained fairly constant [17]. Although it was known that the mean age of onset of T2D in Asian Indians occurs at least a decade or two earlier than in Europeans [17], there is now increasing evidence that it is occurring more frequently among children and adolescents [18].

The problem with the earlier age of onset is that it increases the period of exposure to hyperglycemia, and thus increases the risk of developing long-term diabetic complications by the time they reach early adulthood or midlife. In a study of children and adolescents with a duration of T2D more than 15 years, 81.5% were found to have retinopathy




FIGURE 1. The rising trend of diabetes in India in last two decades. Data from [4–11].

and 34.4% to have nephropathy [19], suggesting a higher rate of complications among young T2D. Similarly, a number of studies suggested that early onset T2D may have a more aggressive course than older onset T2D [20,21] or even T1D [22].

Factors contributing to the rise in prevalence of diabetes in India

Socioeconomic conditions are progressively getting better in India during the last two decades, and India has transitioned from lower economic status to middle economic status. The prevalence of most communicable diseases is decreasing, whereas there is a rise in noncommunicable diseases such as diabetes, hypertension, and heart disease. At present, India faces an epidemiologic transition with a 'double burden' of communicable and noncommunicable diseases [23].

The increased availability of high calorie refined and processed foods, introduction of Western diets, and changes from traditionally healthier food habits to 'junk' food higher in fat and sugar has contributed to a rapid rise in obesity and T2D [2,24]. Recent studies have shown a high intake of refined carbohydrates such as white rice and refined wheat is associated with the high prevalence of diabetes in India [25]. Refined cereal intake was also linked to metabolic syndrome [26]. Conversely, the substitution of white rice with brown rice led to decreased blood glucose and insulin levels [27]. Physical inactivity has also been shown to be a powerful risk factor for diabetes [28]. Psychosocial factors such as depression may also play a role [29]. Another study from our group showed that five factors – unhealthy diet, obesity, physical inactivity, serum triglycerides, and low HDL cholesterol – explain 80.7% of the population attributable risk for diabetes in this population [30].

Several evidences suggest there is a strong genetic component to T2D among Asian Indians: first, a recent study showed the highest incidence of T2D among Asian Indians compared with any ethnic group except Pima Indians, but the latter are much more obese [31]; second, the prevalence of T2D is higher in migrant Indians compared with native populations (e.g., Europeans, Chinese, or Africans) even after an adjustment to age, sex, and BMI [32–35]; third, 36% of Europeans had first-degree relatives with T2D compared with 45% of Asian Indians [36]; fourth, nearly 60% of offspring with two T2D parents had diabetes or impaired glucose tolerance higher than reported in Caucasian populations [37]; and fifth, there is a strong familial aggregation of quantitative traits

that are typically associated with T2D [38]. Recent genome-wide association studies have shown that there are some unique genes associated with T2D in Indians [39–41]. As the genes have not changed drastically during the last two decades, whereas the prevalence of diabetes has increased over 10-fold, it is likely that environmental factors are playing a major role in the escalation of the diabetes prevalence in India.

Another hypothesis that explains the increasing prevalence of T2D in India is the 'thrifty genotype hypothesis' [42]. This hypothesis proposed that some genes are selected over previous millennia to allow survival in times of famine by efficiently storing all available energy during times of feast [42]. With the change in socioeconomic status and the continuous availability of calorie dense foods, there is an increase in obesity and T2D in India because of fat accumulation from the role of these genes. This hypothesis also explains the rising prevalence of T2D in American Indians [43].

Studies show that migration from rural areas to urban slums in India led to high rates of obesity, glucose intolerance, and dyslipidemia as a result of a sedentary lifestyle, stress, and higher intake of refined food [44,45].

Low birth weights have been shown to be associated with T2D in several populations [46–48] including Indians [49]. Other factors, such as exposure to industrial chemicals, smoking, sleep deprivation, and/or depression/stress, are a result of modern lifestyles that might also contribute to an increased prevalence of T2D [50–52]. These factors have, however, not been studied well in the Indian population, and more studies on these factors are urgently needed. Various factors linked to rise in the prevalence of diabetes are summarized in Table 1.

Is type 2 diabetes different in India?

Several studies from migrant Indians and among Indians in India have suggested that there is a specific phenotype that makes Indians distinct from Caucasians (or indeed other ethnic groups) and this is referred to as the 'Asian Indian phenotype' [53,54] [Fig. 2].

The Asian Indian phenotype features are as follows.

Onset of diabetes at a young age

In the recent ICMR-INDIAB national diabetes survey, over half of the subjects had an onset of diabetes below 50 years of age [11] that is higher than among people of European origin [55,56].

Table 1. Factors contributing to the rise of diabetes prevalence among Indians

| Factors (reference) | |
|-----------------------------------|---|
| Diet [24–26,27 ^a] | Higher intake of refined carbohydrates and fats Higher intake of 'junk' food Increase in calorie consumption |
| Physical inactivity [28,30] | Urbanization and sedentary life style Abdominal obesity Hypertension Dyslipidemia |
| Genetics [31 ^a ,38–43] | Higher prevalence compared with other ethnic groups Strong familial aggregation Unique genes associated with T2D in Indians Abdominal obesity, greater insulin resistance Thrifty genotype hypothesis |
| Industrialization [44,45] | Higher stress, sleep deprivation and higher depression among Urban Indians Migration from rural to urban area increases the risk for obesity and metabolic syndrome |
| Low birth weight [46,49] | Studies have shown link between low birth weight and later development of T2D in Indians |
| Others [50–52] | Industrial chemicals Smoking/tobacco Alcohol consumption |

A study from Canada reported that the median age at diagnosis was lowest among South Asians compared with other ethnic groups including people of other Asian countries [57].

Central (abdominal) obesity

Studies have shown that Asian Indians have larger waist circumferences and abdominal obesity compared with Caucasians despite similar age, BMI, and




FIGURE 2. Asian Indian phenotype.

total body fat content [58,59]. Such differences in body fat composition are even seen in childhood and adolescence. Indeed, studies have shown that South Asian neonates have significantly more body fat even on the day of birth compared with European children, despite having lower birth weights [60,61].

Greater insulin resistance

Studies have shown that Asian Indians with and without diabetes are more insulin resistant than Caucasians, even at younger ages and comparative levels of BMI and regardless of total body fat [62]. Abdominal obesity might be responsible for insulin resistant and development of diabetes at younger age [63].

Early decline in beta cell function

A study examining both insulin resistant and beta cell function in a group of East Asians, South Asians, Africans, and Caucasians found that despite being matched for lifestyle factors and BMI, the prevalence of insulin resistance in South Asian men was three to four-fold higher than in other ethnic groups [64]. It was also observed that beta cell responsiveness in South Asians was not sufficient to compensate for the degree of insulin resistance [64]. In addition, studies have shown that Asian Indians lose beta cell function very early in the natural history of T2D, even at the stage of prediabetes [65,66]. A prospective study of Asian Indians living in South Africa showed that participants with impaired glucose tolerance exhibited delayed insulin responses indicating early beta cell dysfunction as an underlying pathophysiological abnormality in development of T2D [67].

Asian Indian genotype

Though some T2D genes are common to Asian Indians and European populations [68–70], factors that mediate genetic effects (allele frequencies or polymorphisms) may differ between Asian Indians and Europeans. For example, the fat mass and obesity associated (FTO) gene is associated with T2D in both Europeans and South Asians; however, the association is entirely mediated by BMI in Europeans but not in Asians [69–70]. Moreover, there are unique genes associated with T2D in Indians [39–41,68]. Studies have also found that genetic polymorphisms in various genes are associated with a higher rate of diabetes complications in Indians [71–73]. Since the Indian population is genetically highly diverse, the findings from different studies should be interpreted while taking into account the geographical area and the ethnic group [74].

Increased cardiovascular risk

In an analysis of age-standardized coronary heart disease (CHD) mortality in Canada over a 15-year period, South Asians had the highest CHD mortality compared with individuals of Chinese and European descent [75]. In addition, South Asian men are prone to developing CHD at a younger stage, often before the age of 40 years [76]. The higher risk for CHD events can be attributed to the unique dyslipidemia pattern with low HDL-C, high serum triglycerides, and increased small dense LDL-C, in addition to other factors such as abnormal adipocytokines due to abnormal visceral fat accumulation. The higher levels of serum leptin and C-reactive protein and lower levels of serum adiponectin have also been linked to the metabolic syndrome and high risk for cardiovascular events in Indians [77–79].

Differences in risk of microvascular complications

There are very few population-based studies on complications of diabetes in India. Studies among Asian Indians have shown lower prevalence of diabetic retinopathy [80–83] and higher prevalence of nephropathy [84–87]. Even given the slightly lower prevalence of microvascular complications, the sheer numbers of diabetic patients in India mean that millions are at risk of developing these complications. Shah *et al.* [88] noted that half of the T2D lack the knowledge regarding diabetes and its treatment that might potentially increase the prevalence of complications.

Future directions

Given the enormity of the noncommunicable diseases burden and its economic impact, the Government of India launched the National Programme for Prevention and Control of Diabetes, Cardiovascular Disease, Stroke and Cancer [89]. The objective was to prevent the rise in communicable disease and complications by early diagnosis and treatment.

Despite much progress in characterizing T2D in Indians, many questions remain about why early beta cell dysfunction occurs in Asian Indians. This should include sophisticated in-depth research such as the euglycemic clamp studies. More mechanistic studies should be undertaken to better characterize the 'Asian Indian phenotype' with particular reference to studies on visceral fat and muscle. Finally, it would be worthwhile examining whether lifestyle changes can override genetic factors, and thus alter the Asian Indian phenotype.

CONCLUSION

The prevalence of type 2 diabetes is rising at an alarming rate in India and now exceeds the rates seen in migrant Indians in the USA, indicating a complete reversal of the scenario seen two or three decades ago. The phenotype of an Asian Indian with type 2 diabetes is different than Caucasians and characterized by higher body fat percentage, higher waist circumference or waist-to-hip ratio representing central obesity, rapid decline in beta cell function, and a greater degree of insulin resistance with higher risk for cardiovascular disease. Clearly, more studies are needed to characterize this unique 'Asian Indian Phenotype'. Finally, prevention and control of diabetes should become a public health priority in India. The time for action is NOW!

Acknowledgements

The authors acknowledge the help of Dawn White and Anne Caulfield at Barbara Davis Center for Diabetes for editing this manuscript.

Financial support and sponsorship

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Das AK, Shah S. History of diabetes: from ants to analogs. *J Assoc Physicians India* 2011; 59 (Suppl):6–7.
2. Weaver LJ, Narayan KM. Reconsidering the history of type 2 diabetes in India: emerging or re-emerging disease? *Natl Med J India* 2008; 21:288–291.
3. Smith B. Eaters, food and social hierarchy in ancient India: a dietary guide to a revolution of values. *J Am Acad Rel* 1990; 58:177–205.
4. Ahuja MMS. Epidemiological studies on diabetes mellitus in India. In: Ahuja MMS, editor. *Epidemiology of diabetes in developing countries*. New Delhi: Interprint; 1979. pp. 29–38.
5. Ramachandran A, Jali MV, Mohan V, et al. High prevalence of diabetes in an urban population in South India. *Br Med J* 1988; 297:587–590.
6. Rao PV, Ushabala P, Seshiah V, et al. The Eluru survey: prevalence of known diabetes in a rural Indian population. *Diabetes Res Clin Pract* 1989; 7:29–31.
7. Ahuja MMS. Recent contributions to the epidemiology of diabetes mellitus in India. *Int J Diabetes Dev Ctries* 1991; 11:5–9.
8. Ramachandran A, Snehalatha C, Kapur A, et al. Diabetes Epidemiology Study Group in India (DESI). High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. *Diabetologia* 2001; 44:1094–1101.
9. Sadikot SM, Nigam A, Das S, et al. Diabetes India. The burden of diabetes and impaired fasting glucose in India using the ADA 1997 criteria: Prevalence of Diabetes in India Study (PODIS). *Diabetes Res Clin Pract* 2004; 66:293–300.
10. Ajay VS, Prabhakaran D, Jeemon P, et al. Prevalence and determinants of diabetes mellitus in the Indian industrial population. *Diabet Med* 2008; 25:1187–1194.
11. Anjana RM, Pradeepa R, Deepa M, et al. On behalf of the ICMR-INDIAB Collaborative Study Group. Prevalence of diabetes and prediabetes (impaired fasting glucose and/or impaired glucose tolerance) in urban and rural India: phase I results of the Indian Council of Medical Research-India DIABetes (ICMR-INDIAB) study. *Diabetologia* 2011; 54:3022–3027.

12. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes Care* 1998; 21:1414–1431.
13. Wild S, Roglic G, Green A, et al. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004; 27:1047–1053.
14. International Diabetes Federation. The diabetes atlas, 6th ed. Poster update. 2014. Available at http://www.idf.org/sites/default/files/Atlas-poster-2014_EN.pdf. [Accessed 2 February 2015]
15. Zargar AH, Wani AA, Laway BA, et al. Prevalence of diabetes mellitus and other abnormalities of glucose tolerance in young adults aged 20–40 years in North India (Kashmir Valley). *Diabetes Res Clin Pract* 2008; 82:276–281.
16. Mohan V, Revale J, Deepa R. Type 2 diabetes in Asian Indian youth. *Pediatr Diabetes* 2007; 8:28–34.
17. Amutha A, Datta M, Unnikrishnan IR, et al. Clinical profile of diabetes in the young seen between 1992 and 2009 at a specialist diabetes centre in south India. *Prim Care Diabetes* 2011; 5:223–229.
18. Mohan V, Deepa M, Deepa R, et al. Secular trends in the prevalence of diabetes and impaired glucose tolerance in urban south India – the Chennai Urban Rural Epidemiology Study (CURES-17). *Diabetologia* 2006; 49:1175–1178.
19. Amutha A, Datta M, Unnikrishnan R, et al. Clinical profile and complications of childhood- and adolescent-onset type 2 diabetes seen at a diabetes centre in south India. *Diabetes Technol Ther* 2012; 14:497–504.
20. Dart A, Martens P, Rigatto C, et al. Earlier onset of complications in youth with type 2 diabetes. *Diabetes Care* 2014; 37:436–443.
21. TODAY Study Group. Retinopathy in youth with type 2 diabetes participating in the TODAY clinical trial. *Diabetes Care* 2013; 36:1772–1774.
22. Constantino MI, Molyneaux L, Limacher-Gisler F, et al. Long-term complications and mortality in young-onset diabetes: type 2 diabetes is more hazardous and lethal than type 1 diabetes. *Diabetes Care* 2013; 36:3863–3869.
23. Anjana RM, Ali MK, Pradeepa M, et al. The need for obtaining accurate nationwide estimates of diabetes prevalence in India: rationale for a national study on diabetes. *Indian J Med Res* 2011; 133:369–380.
24. Mohan V, Sandeep S, Deepa R, et al. Epidemiology of type 2 diabetes: Indian scenario. *Indian J Med Res* 2007; 125:217–230.
25. Mohan V, Radhika G, Sathya RM, et al. Dietary carbohydrates, glycaemic load, food groups and newly detected type 2 diabetes among urban Asian Indian population in Chennai, India (Chennai Urban Rural Epidemiology Study 59). *Br J Nutr* 2009; 102:1498–1506.
26. Radhika G, Van Dam RM, Sudha V, et al. Refined grain consumption and the metabolic syndrome in urban Asian Indians (Chennai Urban Rural Epidemiology Study 57). *Metabolism* 2009; 58:675–681.
27. Mohan V, Spiegelman D, Sudha V, et al. Effect of brown rice, white rice, and brown rice with legumes on blood glucose and insulin responses in overweight Asian Indians: a randomized controlled trial. *Diabetes Technol Ther* 2014; 16:317–325.

This study showed that minor change in quality of carbohydrate such as replacing white rice with brown rice improves glycemic out comes in Indians.

28. Mohan V, Shanthi Rani S, Deepa R, et al. Intra urban differences in the prevalence of the metabolic syndrome in southern India: the Chennai Urban Population Study (CUPS-4). *Diabetic Med* 2001; 18:280–287.
29. Poongothai S, Pradeepa R, Ganesan A, Mohan V. Prevalence of depression in a large urban south Indian population: the Chennai Urban Rural Epidemiology Study (Cures – 70). *PLoS One* 2009; 4:e7185.
30. Anjana RM, Sudha V, Nair DH, et al. Diabetes in Asian Indians- how much is preventable? Ten-year follow-up of the Chennai Urban Rural Epidemiology Study (CURES- 142). *Diabetes Res Clin Pract* 2015 [Epub ahead of print].
31. Anjana RM, Shanthirani CS, Deepa M, et al. Incidence of diabetes and prediabetes and predictors of progression among Asian Indians- ten-year follow-up of the Chennai Urban Rural Epidemiology Study (CURES). *Diabetes Care* 2015. [Epub ahead of print]

The first Indian study to estimate incidence of diabetes in Southern part of India and also characterized the factors responsible for progression of diabetes over ten years.

32. Mather HM, Keen H. The Southall Diabetes Survey: prevalence of known diabetes in Asians and Europeans. *Br Med J (Clin Res Ed)* 1985; 291:1081–1084.
33. Omar MA, Seedat MA, Dyer RB, et al. South African Indians show a high prevalence of NIDDM and bimodality in plasma glucose distribution patterns. *Diabetes Care* 1994; 17:70–73.
34. Jenum AK, Holme I, Graff-Iversen S, Birkeland KI. Ethnicity and sex are strong determinants of diabetes in an urban Western society: implications for prevention. *Diabetologia* 2005; 48:435–439.
35. Lee WR. The changing demography of diabetes mellitus in Singapore. *Diab Res Clin Pract* 2000; 50 (Suppl 2):S35–S39.
36. Mohan V, Sharp PS, Aber VR, et al. Family histories of Asian Indian and Europeans noninsulin-dependent diabetic patients. *Pract Diabetes* 1986; 3:254–256.
37. Viswanathan M, Mohan V, Snehalatha C, Ramachandran A. High prevalence of Type 2 (non insulin dependent) diabetes among offspring of conjugal diabetic parents in India. *Diabetologia* 1985; 28:907–910.

38. Mathias RA, Deepa M, Deepa R, *et al.* Heritability of quantitative traits associated with type 2 diabetes mellitus in large multiplex families from south India. *Metabolism* 2009; 58:1439–1445.
39. Kooner JS, Saleheen D, Sim X, *et al.* Genome-wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes susceptibility loci. *Nat Genet* 2011; 43:984–989.
40. Saxena R, Saleheen D, Been LF, *et al.* Genome-wide association study identifies a novel locus contributing to type 2 diabetes susceptibility in Sikhs of Punjabi origin from India. *Diabetes* 2013; 62:1746–1755.
41. Tabassum R, Chauhan G, Dwivedi OP, *et al.* Genome-wide association study for type 2 diabetes in Indians identifies a new susceptibility locus at 2q21. *Diabetes* 2013; 62:977–986.
42. Neel JV, Weder AB, Julius S. Type II diabetes, essential hypertension, and obesity as “syndromes of impaired genetic homeostasis” the “thrifty genotype” hypothesis enter the 21st century. *Perspect Biol Med* 1998; 42:44–74.
43. Burrows NR, Geiss LS, Engelgau MM, Acton KJ. Prevalence of diabetes among Native Americans and Alaska Natives, 1990–1997: an increasing burden. *Diabetes Care* 2000; 23:1786–1790.
44. Misra A, Pandey RM, Devi JR, *et al.* High prevalence of diabetes, obesity and dyslipidaemia in urban slum population in northern India. *Int J Obes Relat Metab Disord* 2001; 25:1722–1729.
45. Deepa M, Anjana RM, Manjula D, *et al.* Convergence of prevalence rates of diabetes and cardio metabolic risk factors in middle and low income groups in urban India: 10-year follow up of the Chennai Urban Population Study. *J Diab Sci Technol* 2011; 5:918–927.
46. Lindsay RS, Dabelea D, Roumain J, *et al.* Type 2 diabetes and low birth weight: the role of paternal inheritance in the association of low birth weight and diabetes. *Diabetes* 2000; 49:445–449.
47. Hypponen E, Smith GD, Power C. Parental diabetes and birth weight of offspring: intergenerational cohort study. *Br Med J* 2003; 326:19–20.
48. Johansson S, Iliadou A, Bergvall N, *et al.* The association between low birth weight and type 2 diabetes: contribution of genetic factors. *Epidemiology* 2008; 19:659–665.
49. Yajnik CS, Coyaji KJ, Joglekar CV, *et al.* Paternal insulin resistance and fetal growth: problem for the ‘fetal insulin’ and the ‘fetal origins’ hypotheses. *Diabetologia* 2001; 44:1197–1198.
50. Roopa M, Deepa M, Indulekha K, Mohan V. Prevalence of sleep abnormalities and their association with metabolic syndrome among Asian Indians: Chennai Urban Rural Epidemiology Study (CURES-67). *J Diab Sci Technol* 2010; 4:1524–1531.
51. Minh NH, Minh TB, Kajiwarana N, *et al.* Contamination by persistent organic pollutants in dumping sites of Asian developing countries: implication of emerging pollution sources. *Arch Environ Contam Toxicol* 2006; 50:474–481.
52. Rimm EB, Chan J, Stampfer MJ. Prospective study of cigarette smoking, alcohol use, and the risk of diabetes in men. *BMJ* 1995; 310:555–559.
53. Deepa R, Sandeep S, Mohan V. Abdominal obesity, visceral fat and type 2 diabetes: “Asian Indian phenotype”. In: Mohan V, Rao GHR, editors. *Type 2 diabetes in South Asians: epidemiology risk factors and prevention*. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd; 2006. pp. 138–152.
54. Unnikrishnan R, Anjana RM, Mohan V. Diabetes in South Asians: is the phenotype different? *Diabetes* 2014; 63:53–55.
55. Center for Disease Control and Prevention. Diabetes data and trends. Available at <http://www.cdc.gov/diabetes/statistics/incidence/>. [Accessed 15 January 2015]
56. Gujral UP, Pradeepa R, Weber MB, *et al.* Type 2 diabetes in South Asians: similarities and differences with white Caucasian and other populations. *Ann N Y Acad Sci* 2013; 1281:51–63.
57. Chiu M, Austin PC, Manuel DG, *et al.* Deriving ethnic-specific BMI cutoff points for assessing diabetes risk. *Diabetes Care* 2011; 34:1741–1748.
58. Raji A, Seely EW, Arky RA, Simonson DC. Body fat distribution and insulin resistance in healthy Asian Indians and Caucasians. *J Clin Endocrinol Metab* 2001; 86:5366–5371.
59. Chandalia M, Abate N, Garg A, *et al.* Relationship between generalized and upper body obesity to insulin resistance in Asian Indian men. *J Clin Endocrinol Metab* 1999; 84:2329–2335.
60. Krishnaveni GV, Hill JC, Veena SR, *et al.* Truncal adiposity is present at birth and in early childhood in South Indian children. *Indian Pediatr* 2005; 42:527–538.
61. Yajnik CS, Fall CH, Coyaji KJ, *et al.* Neonatal anthropometry: the thin-fat Indian baby. The Pune Maternal Nutrition Study. *Int J Obes Relat Metab Disord* 2003; 27:173–180.
62. Mohan V, Sharp PS, Cloke HR, *et al.* Serum immunoreactive insulin responses to a glucose load in Asian Indian and European type 2 (non insulin dependent) diabetic patients and control subjects. *Diabetologia* 1986; 29:235–237.
63. Indulekha K, Anjana RM, Surendar J, Mohan V. Association of visceral and subcutaneous fat with glucose intolerance, insulin resistance, adipocytokines and inflammatory markers in Asian Indians (CURES-113). *Clin Biochem* 2011; 44:281–287.
64. Petersen KF, Dufour S, Feng J, *et al.* Increased prevalence of insulin resistance and nonalcoholic fatty liver disease in Asian-Indian men. *Proc Natl Acad Sci USA* 2006; 103:18273–18277.
65. Staimez LR, Weber MB, Ranjani H, *et al.* Evidence of reduced beta cell function in Asian Indians with mild dysglycemia. *Diabetes Care* 2013; 36:2772–2778.
66. Mohan V, Amutha A, Ranjani H, *et al.* Associations of β -cell function and insulin resistance with youth-onset type 2 diabetes and prediabetes among Asian Indians. *Diabetes Technol Ther* 2013; 15:315–322.
67. Motala AA, Omar MA. Evidence for impaired pancreatic beta cell function in South African Indians with impaired glucose tolerance. *Diabet Med* 1994; 11:437–444.
68. Chauhan G, Spurgeon CJ, Tabassum R, *et al.* Impact of common variants of PPARG, KCNJ11, TCF7L2, SLC30A8, HHEX, CDKN2A, IGF2BP2, and CDKAL1 on the risk of type 2 diabetes in 5164 Indians. *Diabetes* 2010; 59:2068–2074.
69. Yajnik CS, Janipalli CS, Bhaskar S, *et al.* FTO gene variants are strongly associated with type 2 diabetes in South Asian Indians. *Diabetologia* 2009; 52:247–252.
70. Chauhan G, Tabassum R, Mahajan A, *et al.* Common variants of FTO and the risk of obesity and type 2 diabetes in Indians. *J Hum Genet* 2011; 56:720–726.
71. Shah VN, Cheema BS, Sharma R, *et al.* ACAC β gene (rs2268388) and AGTR1 gene (rs5186) polymorphism and the risk of nephropathy in Asian Indian patients with type 2 diabetes. *Mol Cell Biochem* 2013; 372:191–198.
72. Shah VN, Cheema BS, Kohli HS, *et al.* Endothelial nitric oxide synthase gene polymorphism and the risk of diabetic neuropathy in Asian Indian patients with type 2 diabetes. *J Diabetes Metab* 2013; 4:243.
73. Cheema BS, Kohli HS, Sharma R, *et al.* RAS gene polymorphisms and renal responsiveness to RAS inhibition therapy in type 2 diabetic Asian Indians. *J Pharmacogenom Pharmacoproteomics* 2013; 4:114.
74. Holliday EG. Hints of unique genetic effects for type 2 diabetes in India. *Diabetes* 2013; 62:1369–1370.
75. Sheth T, Nair C, Nargundkar M, *et al.* Cardiovascular and cancer mortality among Canadians of European, South Asian and Chinese origin from 1979 to 1993: an analysis of 1.2 million deaths. *Can Med Assoc J* 1999; 161:132–138.
76. Enas EA, Garg A, Davidson MA, *et al.* Coronary heart disease and its risk factors in first-generation immigrant Asian Indians to the United States of America. *Indian Heart J* 1996; 48:343–353.
77. Lau DC, Dhilon B, Yan H, *et al.* Adipokines: molecular links between obesity and atherosclerosis. *Am J Physiol Heart Circ Physiol* 2005; 288:H2031–H2041.
78. Mohan V, Deepa R, Velmurugan K, Premalatha G. Association of C-reactive protein with body fat, diabetes and coronary artery disease in Asian Indians: the Chennai Urban Rural Epidemiology Study (CURES-6). *Diab Med* 2005; 22:863–870.
79. Abate N, Chandalia M, Snell PG, Grundy SM. Adipose tissue metabolites and insulin resistance in nondiabetic Asian Indian men. *J Clin Endocrinol Metab* 2004; 89:2750–2755.
80. Dandona L, Dandona R, Naduvilath TJ, *et al.* Population based assessment of diabetic retinopathy in an urban population in southern India. *Br J Ophthalmol* 1999; 83:937–940.
81. Narendran V, John RK, Raghuram A, *et al.* Diabetic retinopathy among self reported diabetics in southern India: a population based assessment. *Br J Ophthalmol* 2002; 86:1014–1018.
82. Rema M, Premkumar S, Anitha B, *et al.* Prevalence of diabetic retinopathy in urban India: the Chennai Urban Rural Epidemiology Study (CURES) Eye Study - 1. *Invest Ophthalmol Vis Sci* 2005; 46:2328–2333.
83. Raman R, Rani PK, Racheppalle RS, *et al.* Prevalence of diabetic retinopathy in India: Sankara Nethralaya Diabetic Retinopathy Epidemiology and Molecular Genetics Study report 2. *Ophthalmology* 2009; 116:311–318.
84. Samanta A, Burden AC, Feehally J, Walls J. Diabetic renal disease: differences between Asian and white patients. *Br Med J (Clin Res Ed)* 1986; 293:366–367.
85. Mather HM, Chaturvedi N, Fuller JH. Mortality and morbidity from diabetes in South Asians and Europeans: 11-year follow-up of the Southall Diabetes Survey, London, UK. *Diabet Med* 1998; 15:53–59.
86. Chandie Shaw PK, Baboe F, van Es LA, *et al.* South-Asian type 2 diabetic patients have higher incidence and faster progression of renal disease compared with Dutch-European diabetic patients. *Diabetes Care* 2006; 29:1383–1385.
87. Unnikrishnan RI. Diabetic nephropathy in south Asians. In: Mohan V, Rao GHR, editors. *Type 2 diabetes in South Asians: epidemiology, risk factors and prevention*, 1st ed. New Delhi, India: Jaypee Brothers Medical Publishers; 2006. pp. 230–236.
88. Shah VN, Kamdar PK, Shah N. Assessing the knowledge, attitudes and practice of type 2 diabetes among patients of Saurashtra region, Gujarat. *Int J Diabetes Dev Countries* 2009; 29:118–1122.
89. National Programme for Prevention and Control of Diabetes, Cardiovascular Diseases and Stroke (NPDCS). http://www.searo.who.int/india/topics/cardiovascular_diseases/NCD_Resources_Training_module_for_NPDCS_for_health_workers.pdf?ua=1. [Accessed 28 March 2015]